


National Water Conditions

UNITED STATES
Department of the Interior
Geological Survey

CANADA
Department of the Environment
Water Resources Branch

MAY 1986



Both variability and persistence marked streamflow during May as precipitation across the United States varied widely. For example: streamflow went from above normal to below normal in parts of Ontario, Michigan, Wisconsin, Montana, and Arizona; streamflow went from below normal to above normal in parts of Washington, Oregon, California, and Virginia; record high flows for May occurred in the upper midcontinent, record low flows for May occurred in the East; parts of the United States were into their 5th year of above-normal streamflow, the rise of Utah's Great Salt Lake being one symptom of the phenomenon.

On May 30, eight people were killed, one person was missing, and damages were about \$20 million after flooding in the area of Etna, Pennsylvania.

On June 1, 1986, Utah's Great Salt Lake stood at 4,211.80 feet above sea level, 0.20 foot higher than the 1873 record-high of 4,211.6 feet.

The combined flow of the 3 largest rivers in the lower 48 States—Mississippi, St. Lawrence, and Columbia—averaged 1,237,200 cfs during May, 20 percent below median, but 1 percent above last month.

Contents of 71 percent of reporting reservoirs were at or above average for the end of May. Only five reservoirs reported significant declines in contents during May and only two of those, International Amistad in Texas and Hungry Horse in Montana, had below-average contents for the end of May.

About 57 percent of the index stations in the United States and southern Canada had flows in the normal to above-normal range, compared with the 61 percent in these ranges for last month.

STREAMFLOW CONDITIONS DURING MAY 1986

Both variability and persistence marked streamflow during May as precipitation across the United States (see map) varied widely. For example: record low precipitation for May (data from National Weather Service) occurred in Bridgeport, Connecticut (0.34 inches), Marquette, Michigan (0.07 inches), Phoenix, Arizona (0.00 inches), and several other locations, while San Angelo, Texas, had a record-breaking 7.28 inches; streamflow went from above normal to below normal in parts of Ontario, Michigan, Wisconsin, Montana, and Arizona; streamflow went from below normal to above normal in parts of Washington, Oregon, California, and Virginia; record high flows for May occurred in the upper midcontinent, record low flows for May occurred in the East (see table); parts of the United States were into their 5th year of above-normal streamflow (see page 6), the rise of Utah's Great Salt Lake being one symptom of the phenomenon. The four hydrographs on page 4 were chosen to highlight both the variability and persistence of flow conditions across the Nation. Smith River and Marias River have been in three different flow ranges in as many months; Crow River is in its 20th consecutive month of above-normal flow, and set a new record high for May; Contentnea Creek is in its 5th consecutive month of below-normal flow and set a record monthly low for May, as it did in April.

On May 20, the "Upriver Dam" on the Spokane River at the Spokane, Washington, city limits was breached by rising waters after a power failure during a thunderstorm on May 20. The dam gates were closed at the time and the spillway could not convey the 9,000 cubic feet per second (cfs) flow. Peak discharge after the breach was about 16,200 cfs at the Spokane River gaging station 4.5 miles downstream, 10,200 cfs less than the 1985 annual peak at that site.

On May 30, eight people were killed, one person was missing, and damages were about \$20 million after flooding in the area of Etna, Pennsylvania. Little Pine Creek near Etna (drainage area 5.78 square miles) peaked at about 7,400 cfs (recurrence interval greater than 100 years) as a result of 7 inches or more of rain (National Weather Service bucket survey) in the headwaters area of the creek, over three times the previous record peak discharge (2,040 cfs in 1974).

On June 1, 1986, Utah's Great Salt Lake stood at 4,211.80 feet above sea level, 0.20 foot higher than the 1873 record-high of 4,211.6 feet. Records of lake level began in 1847. The lake rose only 0.15 foot from May 15 to June 1, and the 0.5-foot rise recorded during May was the smallest monthly rise since January 1986. The National Weather Service revised the peak forecast to 4,212 feet based on the June 1 reading. (A dike

breached on June 8 when the lake stood at 4,211.85 feet, and the combination of evaporation and flooding of the previously protected area dropped the lake elevation to 4,211.45 feet as of June 10.)


The combined flow of the 3 largest rivers in the lower 48 States—Mississippi, St. Lawrence, and Columbia—averaged 1,237,200 cfs during May, 20 percent below median, but 1 percent above last month. Discharges increased on the St. Lawrence River (4 percent) and the Columbia River (25 percent) from last month, somewhat more than offsetting the 9 percent decrease in flow of the Mississippi River. The combined flow of the three rivers and the individual flows of both the Mississippi and Columbia were in the below-normal range, but the flow of the St. Lawrence was 19 percent above median, in the above-normal range for the 16th consecutive month, and the 2nd highest May flow in 127 years of record. These three rivers account for runoff from more than half the conterminous United States and provide a useful check on the status of the Nation's surface-water resources.

Contents of 71 percent of reporting reservoirs were at or above average for the end of May. Only five reservoirs reported significant declines in contents during May and only two of those, International Amistad in Texas and Hungry Horse in Montana, had below-average contents for the end of May. In sharp contrast, 13 reservoirs reported both significant increases in contents during May and above-average contents for the end of May. Contents of Belle Fourche in South Dakota went from a below-average 62 percent of capacity at the end of April to an above-average 86 percent of capacity at the end of May. Contents of the Mississippi River headwater system reservoirs, which more than doubled last month to an above-average 44 percent capacity, again increased significantly to 53 percent of capacity, well above average for the end of May.

Streamflow generally decreased seasonally in Hawaii, Arizona, South Dakota, Minnesota, Missouri, Arkansas, Wisconsin, Illinois, Michigan, Ohio, Pennsylvania, New York, New England, New Brunswick, Nova Scotia, New Jersey, Maryland, Virginia, the Carolinas, Georgia, Florida, Alabama, and Tennessee, and decreased variably in Ontario and Nebraska. Flows changed variably in California, New Mexico, North Dakota, and Indiana, but remained unchanged in Nevada. Flows generally increased in the rest of southern Canada and the United States: seasonally in Quebec, Alberta, British Columbia, Iowa, Kansas, Oklahoma, Texas, Montana, Idaho, Wyoming, Colorado, Utah, and Alaska; contraseasonally in Saskatchewan, Oregon, Louisiana, Mississippi, Kentucky, and West Virginia; variably in Puerto Rico. About 57 percent of the index stations in the United States and

CONTENTS

	Page
Streamflow (map).....	1
Streamflow conditions.....	2
Total precipitation (map).....	3
Palmer Drought Index maps.....	5
Wet trend in the United States.....	6
Ground-water conditions.....	8
Usable contents of selected reservoirs.....	10
Usable contents of selected reservoirs (graphs).....	11
Flow of large rivers.....	12
Dissolved solids and water temperatures at downstream sites on six large rivers.....	13
Precipitation forecast for June through August 1986.....	13
Streamflow during May, 1981-86.....	14
Explanation of data.....	15



NEW EXTREMES DURING MAY 1986 AT STREAMFLOW INDEX STATIONS

Station number	Stream and place of determination	Drainage area (square miles)	Years of record	Previous May extremes (period of record)		May 1986			
				Monthly mean in cfs (year)	Daily mean in cfs (year)	Monthly mean in cfs	Percent of median	Daily mean in cfs	Day
LOW FLOWS									
01E0001	St. Mary's River at Stillwater, Nova Scotia, Canada.	52,311	71	678 (1960)	187 (1960)	597	35	299	23
01FB001	Northeast Margaree River at Margaree Valley, Nova Scotia, Canada.	142	70	463 (1983)	286 (1983)	252	15	139	31
01309500	Massapequa Creek at Massapequa, NY.	38	83	5.01 (1981)	2.80 (1966)	3.07	26	2.30	23
02091500	Contentnea Creek at Hookerton, NC.	729	58	99.1 (1981)	58 (1955)	87	18	44	13
02118000	South Yadkin River near Mocksville, NC.	306	48	146 (1941)	100 (1941)	124	37	107	10
02131000	Pee Dee River at Peedee, SC.	8,830	48	3,083 (1981)	1,200 (1981)	2,516	33
02226000	Altamaha River at Doctortown, GA.	13,600	55	4,326 (1981)	2,820 (1981)	2,960	24	2,580	29
02256500	Fisheating Creek at Palmdale, FL.	311	55	0.0 (1985)	0.0 (1932)	0.0	0	0.0	(*)
02317500	Alapaha River at Statenville, GA.	1,400	51	86.4 (1968)	38 (1938)	78	12	54	31
02358000	Apalachicola River at Chattahoochee, FL.	17,300	57	9,840 (1941)	6,990 (1941)	9,190	46
HIGH FLOWS									
05062000	Buffalo River near Dilworth, MN.	1,040	55	906 (1962)	3,100 (1962)	923	442	2,000	14
05280000	Crow River at Rockford, MN.	2,520	61	4,564 (1975)	11,000 (1906)	6,020	492	8,790	5
05288500	Mississippi River near Anoka, MN.	19,100	55	38,490 (1950)	58,800 (1975)	39,965	266	49,700	16
05330000	Minnesota River near Jordan, MN.	16,200	52	20,630 (1944)	35,100 (1960)	22,248	404	35,800	5
05331000	Mississippi River at St. Paul, MN.	36,800	88	48,460 (1975)	78,100 (1975)	64,232	290	83,900	5
06485500	Big Sioux River at Akron, IA.	9,030	58	6,930 (1984)	11,100 (1984)	7,830	913	16,300	1

*Occurred more than once.

southern Canada had flows in the normal to above-normal range, compared with the 61 percent in these ranges for last month. About 68 percent of the index stations in the United States east of the Mississippi River had flows in the below-normal range compared with about 71 percent in that range for last month.

Below-normal streamflow persisted in Rhode Island, Kentucky, the Carolinas, Georgia, Alabama, and also in parts of Maine, New Hampshire, Vermont, Massachusetts, Connecticut, New York, Pennsylvania, Delaware, Maryland, Virginia, West Virginia, Ohio, Illinois, Tennessee, Louisiana, Mississippi, Florida, and Washington. Flows moved into the below-normal range in parts of Alaska, British Columbia, Washington, Idaho, Montana, California, Arizona, Ontario, Michigan, Wisconsin, Quebec, Nova Scotia, Maine, New Hampshire, Vermont, New York, New Jersey, Pennsylvania,





Maryland, Delaware, and Florida, and also in all of New Brunswick.



Above-normal streamflow persisted in parts of Hawaii, Quebec, Oklahoma, Ontario, Manitoba, Minnesota, Wisconsin, Illinois, Iowa, Missouri, the Dakotas, Nebraska, Montana, Wyoming, Colorado, New Mexico, Idaho, Utah, Nevada, and California. Flows moved into the above-normal range in Puerto Rico, and also in parts of Hawaii, Washington, Oregon, California, Montana, the Dakotas, Manitoba, Minnesota, Wisconsin, Illinois, Iowa, Missouri, Virginia, and Quebec.

Flood stages, as designated by the National Weather Service, were exceeded on many rivers and small streams in Washington, Idaho, Nevada, California, Montana, Colorado, New Mexico, the Dakotas, Nebraska, Kansas, Oklahoma, Texas, Minnesota, Iowa, Missouri, Arkansas, Louisiana, Mississippi, Wisconsin, Illinois, Indiana, Kentucky, New York, New Jersey, and Florida.

SURFACE WATER — MONTHLY MEAN DISCHARGE IN KEY STREAMS

Unshaded area indicates range between highest and lowest record for the month. Dashed line indicates median of monthly values for reference period, 1951-80. Heavy line indicates mean for current period.





(From *Weekly Weather and Crop Bulletin* prepared and published by the NOAA/USDA Joint Agricultural Weather Facility)

WET TREND IN THE UNITED STATES


The rise of the Great Salt Lake to an elevation higher than that observed by the Mormon settlers in 1873 (they arrived in Utah in 1847) is just one symptom of a wet trend affecting two large areas (and several small areas) of the Nation. The first area includes parts of Utah and adjacent States and extends into Oregon; the second area includes parts of the Mississippi River basin and the western Great Lakes/St. Lawrence River basin, as can be seen on the map showing composite annual (calendar year) streamflow from 1982 through 1985. (The composite map was produced by overlaying the four annual maps and appropriately shading those areas with above-normal flows in all 4 years and above- or below-normal flows in 3 out of 4 years.) In contrast, parts of Kansas, Montana, and Hawaii have had 3 of 4 years in the below-normal flow range, but no part of the Nation has had below-normal flows for 4 consecutive years.


Two excellent measures of wetness in the western area are the elevation of Great Salt Lake, for which records (or computations) go back to 1847, and the flow of the Colorado River near Cisco, Utah, for which records go back to 1895. The 12.15-foot rise of the Great Salt Lake from 4,199.65 feet on September 15, 1982, to 4,211.80 feet on June 1, 1986, exceeded in both magnitude and rapidity the 12-foot rise from 4,199.6 feet in 1861 to 4,211.6 feet in 1873. (The data presented are for the south arm of the lake only.) The seasonal rises of the lake in 1982-83, 1983-84, and 1985-86 (through June 1) rank 1, 2, 3 in elevation, and 2, 1, 3 in volume, respectively, for the period 1847-1986. Total increase in volume from the 1982 seasonal low to June 1, 1986, was 9,486,140 acre-feet (3,092 billion gallons)—enough water to cover Rhode Island (area 1,214 square miles) with

12.21 feet of water. The hydrograph of lake elevations from the 1980 seasonal low to June 1, 1986, shows lake fluctuations in detail and illustrates the cumulative effect of excess precipitation in the area. The hydrograph of the Colorado River near Cisco, Utah (which is the major runoff-producing area of the Colorado River basin), shows the immediate effect of excess precipitation at the core of the wet area. Flow at this index station has been in the above-normal range for 45 of the last 46 months, 37 of them consecutive. What the hydrograph does not show is that June is the only month for which a monthly mean discharge observed during 1983-86 is not within the three highest of record and that 23 of the 36 slots for the three highest flows of each individual month are occupied by flows which occurred during 1983-86 (including all of those for November through March)—evidence of an unparalleled period of above-normal flow in the area.



On the fringes of the second large area on the other side of the Continental Divide, Lake Superior and Lake Erie have been at record high levels for several months, flow of the St. Lawrence River during February was a record high for the month with flows during March, April, and May 1986 the second highest ever recorded for those months, and both Lake Huron and Lake Ontario are at well above normal levels, as can be seen on the hydrographs. Flow of the St. Lawrence River has been in the above-normal range for 25 of the last 31 months, 16 of them consecutive. Crow River at Rockford, Minnesota, (hydrograph on page 4), in the core of this area of above-normal flows, has been in the above-normal range for 36 of the last 44 months, the last 20 of them consecutive.

COMPOSITE STREAMFLOW FOR CALENDAR YEARS 1982 THROUGH 1985











Great Salt Lake, UT, Stage 1st and 15th of Month



Unshaded area indicates range between highest and lowest record for the month. Dashed line indicates median of monthly values for reference period, 1951-80. Heavy line indicates mean for current period. (Great Lakes elevations from National Weather Service.)



LOCATION OF SELECTED GAGES ON THE GREAT LAKES AND THE ST. LAWRENCE RIVER




GROUND-WATER CONDITIONS DURING MAY 1986

Ground-water levels declined seasonally in most of the region. (See map.) The principal exceptions were rises in water levels in parts of western New York and Pennsylvania. Ground-water levels near the end of the month were above average in parts of Maine and western New York State. Below-average levels persisted in much of central New England, and levels were also below-average in much of Maryland, Delaware, and southern New Jersey.

In the Southeastern States, ground-water levels showed mixed trends in West Virginia, Kentucky, Arkansas, and Louisiana. Water levels declined in Virginia and North Carolina, and in most wells in Mississippi and Georgia. Water levels were mixed with respect to average in West Virginia, Louisiana, and Florida, and were below average in Kentucky, Virginia, North Carolina, Arkansas, and Mississippi. New low ground-water levels for May were reached in Tennessee, Mississippi, and Georgia, and the water level in the key well in the Savannah area in the coastal plain of Georgia declined to a new all-time low in 30 years of record.

In the central and western Great Lakes States, ground-water levels rose in Iowa, showed mixed trends in Minnesota, and declined in Michigan and Ohio. Water





levels were above average in Minnesota and Michigan, mostly above average in Iowa, and generally below average in Indiana and Ohio.



Map showing ground-water storage near end of May and change in ground-water storage from end of April to end of May.

MONTH-END GROUND-WATER LEVELS IN KEY WELLS

Unshaded area indicates range between highest and lowest record for the month. Dashed line indicates average of monthly levels in previous years. Heavy line indicates level for current period.



In the Western States, ground-water levels rose in Arizona and in most of the key wells in Idaho, and declined in Nevada. Trends were mixed in other Western States. Water levels were above average in North Dakota and Nebraska, and were mixed with respect to average in other States. New high water levels for May were recorded in the Steptoe Valley well in Nevada, and in the

Berrendo-Smith well in New Mexico, despite slight net declines during the month. The level in the key well in the Blanding area in Utah rose, also reaching a new high level for May. Levels declined to new May low levels in the key well at the Kansas Agricultural Experiment Station in Colby, and in the El Paso well in Texas.

Provisional data; subject to revision

WATER LEVELS IN KEY OBSERVATION WELLS IN SOME REPRESENTATIVE AQUIFERS IN THE CONTERMINOUS UNITED STATES—MAY 1986

Aquifer and Location	Water level in feet with reference to land-surface datum	Departure from average in feet	Net change in water level in feet since:		Year records began	Remarks
			Last month	Last year		
Glacial drift at Hanska, south-central Minnesota.	-3.86	+1.67	-1.19	-0.88	1942	
Glacial drift at Roscommon in north-central part of Lower Peninsula, Michigan.	-3.54	+0.39	-0.23	-0.46	1935	
Glacial drift at Marion, Iowa.....	-2.28	+1.41	+0.52	+1.47	1941	
Glacial drift at Princeton in northwestern Illinois.	-6.44	+1.49	+1.54	+2.08	1943	
Petersburg Granite, southeastern Piedmont near Fall Zone, Colonial Heights, Virginia.	-15.09	-0.25	-0.45	+1.23	1939	
Glacial outwash sand and gravel, Louisville, Kentucky (U.S. well no. 2).	-17.50	+7.42	-0.02	-1.03	1946	
500-foot sand aquifer near Memphis, Tennessee (U.S. well no. 2).	-104.73	-15.91	-0.06	-1.07	1941	May low.
Granite in eastern Piedmont Province, Chapel Hill, North Carolina (U.S. well no. 5).	-42.29	-1.11	-0.32	-1.84	1931	
Sparta Sand in Pine Bluff industrial area, Arkansas.	-219.15	-12.51	-1.05	+3.35	1958	
Eutaw Formation in the City of Montgomery, Alabama (U.S. well no. 4).	-25.0	-4.6	-0.2	-3.2	1952	
Limestone aquifer on Cocksbur Island, Savannah area, Georgia (U.S. well no. 6).	-35.18	-8.84	-2.09	+0.94	1956	Alltime low.
Sand and gravel in Puget Trough, Tacoma, Washington.	-99.78	+9.60	+0.32	+8.50	1952	
Pleistocene glacial outwash gravel, North Pole, northern Idaho (U.S. well no. 3).	-461.5	-1.0	+0.2	-2.5	1929	
Snake River Group: Snake River Plain Aquifer, at Eden, Idaho (U.S. well no. 4).	-123.1	-2.3	+1.2	+0.9	1957	
Alluvial valley fill in Flowell area, Millard County, Utah (U.S. well no. 9).	-9.08	+28.45	-0.29	-1.81	1929	
Alluvial sand and gravel, Platte River Valley, Ashland, Nebraska (U.S. well no. 6).	-3.20	+0.96	+0.34	-0.57	1935	
Alluvial Valley fill in Steptoe Valley, Nevada.	-7.07	+5.49	-0.33	+0.59	1950	May high.
Pleistocene terrace deposits in Kansas River valley, at Lawrence, northeastern Kansas.	-17.08	+3.59	+1.14	+1.11	1953	
Alluvium and Paso Robles clay, sand, and gravel, Santa Maria Valley, California	-127.71	+13.30	-15.75	-28.02	1957	
Valley fill, Elfrida area, Douglas, Arizona (U.S. well no. 15).	-104.4	-23.9	+0.4	+0.9	1951	
Hueco bolson, El Paso area, Texas.....	-266.18	-18.39	-1.38	-1.67	1965	May low.
Evangelina aquifer, Houston area, Texas.....	-305.60	-9.85	+1.90	-4.18	1965	








USABLE CONTENTS OF SELECTED RESERVOIRS NEAR END OF MAY 1986

[Contents are expressed in percent of reservoir capacity. The usable storage capacity of each reservoir is shown in the column headed "Normal maximum."]

Principal uses: F-Flood control I-Irrigation M-Municipal P-Power R-Recreation W-Industrial	Reservoir				Percent of normal maximum	Principal uses: F-Flood control I-Irrigation M-Municipal P-Power R-Recreation W-Industrial	Reservoir				Percent of normal maximum	Normal maximum (acre-feet)	
	End of May 1986	End of May 1985	Average for end of May	End of Apr. 1986			End of May 1986	End of May 1985	Average for end of May	End of Apr. 1986			
NOVA SCOTIA													
Rossignol, Mulgrave, Falls Lake, St. Margaret's Bay, Black, and Ponhook Reservoirs (P)	57	47	78	56	226,300	NEBRASKA							1,948,000
QUEBEC													
Allard (P)	83	85	88	76	280,600	OKLAHOMA							2,378,000
Gouin (P)	75	82	65	57	6,954,000	Eufaula (FRP)	113	98	97	99	661,000		
MAINE													
Seven reservoir systems (MP)	89	75	90	85	4,107,000	Keystone (FPR)	107	100	107	85	628,200		
NEW HAMPSHIRE													
First Connecticut Lake (P)	90	97	87	67	76,450	Tenkiller Ferry (FPR)	109	105	103	109	628,200		
Lake Francis (FPR)	72	80	83	62	99,310	Lake Altus (FIMR)	31	25	66	30	133,000		
Lake Winnepesaukee (FR)	102	82	102	94	165,700	Lake O'The Cherokees (FPR)	94	97	94	96	1,492,000		
VERMONT													
Harriman (P)	91	94	87	86	116,200	OKLAHOMA-TEXAS							2,722,000
Somerset (P)	86	77	86	80	57,390	Lake Texoma (FIMPRW)	100	98	103	96			
MASSACHUSETTS													
Cobble Mountain and Borden Brook (MP)	82	65	90	84	77,920	TEXAS							
NEW YORK													
Great Sacandaga Lake (FPR)	100	92	97	90	786,700	Bridgeport (IMW)	90	82	54	79	386,400		
Indian Lake (FMP)	103	92	103	96	103,300	Canyon (FMR)	99	87	81	96	385,600		
New York City reservoir system (MW)	100	61	100	99	1,680,000	International Amistad (FIMPW)	42	63	81	54	3,497,000		
NEW JERSEY													
Wanaque (M)	97	68	95	101	85,100	International Falcon (FIMPW)	28	36	65	24	2,668,000		
PENNSYLVANIA													
Allegheny (FPR)	51	46	47	52	1,180,000	Livingston (IMW)	105	102	91	101	1,788,000		
Pymatung (FMR)	101	95	99	96	188,000	Potomac Kingdom (IMPRW)	92	97	98	86	570,200		
Raystown Lake (FR)	68	68	61	68	761,900	Red Bluff (P)	19	27	26	21	307,000		
Lake Wallenpaupack (FR)	80	90	79	81	157,800	Toledo Bend (P)	97	101	93	93	4,472,000		
MARYLAND													
Baltimore municipal system (M)	84	94	95	86	261,900	Twin Buttes (FIM)	11	12	31	11	177,800		
NORTH CAROLINA													
Bridgewater (Lake James) (P)	92	92	92	90	288,800	Lake Kemp (IMW)	93	89	89	89	268,000		
Narrowes (Bald Lake) (P)	89	94	99	90	128,900	Lake Meredith (FWM)	28	35	36	29	796,900		
High Rock Lake (P)	73	78	83	67	234,800	Lake Travis (FIMPRW)	94	93	83	93	1,144,000		
SOUTH CAROLINA													
Lake Murray (P)	89	87	84	89	1,614,000	MONTANA							
Lakes Marion and Moultrie (P)	80	82	80	80	1,862,000	Canyon Ferry (FIMPR)	81	82	80	75	2,043,000		
SOUTH CAROLINA-GEORGIA													
Clark Hill (FP)	56	67	76	57	1,730,000	Fort Peck (FPR)	80	81	86	76	18,910,000		
GEORGIA													
Burton (PR)	100	96	94	92	104,000	Hungry Horse (FIPR)	65	87	74	79	3,451,000		
Sinclair (MPR)	87	88	93	89	214,000	WASHINGTON							
Lake Sidney Lanier (FMPR)	49	64	66	50	1,686,000	Ross (PR)	71	59	57	51	1,052,000		
ALABAMA													
Lake Martin (P)	91	99	95	92	1,375,000	Franklin D. Roosevelt Lake (IP)	63	33	73	55	5,022,000		
TENNESSEE VALLEY													
Clinch Projects: Norris and Melton Hill Lakes (FPR)	52	55	66	51	2,293,000	Lake Chelan (PR)	90	63	73	57	676,100		
Douglas Lake (FPR)	41	45	72	36	1,394,000	Lake Cushman (PR)	102	72	95	98	359,500		
Hiwassee Projects: Chatuge, Nottely, Hiwassee, Apalachia, Blue Ridge, Ocoee 3, and Parksville Lakes (FPR)	62	67	83	60	1,012,000	Lake Merwin (P)	104	107	104	103	245,600		
Holston Projects: South Holston, Watauga, Boone, Fort Patrick Henry, and Cherokee Lakes (FPR)	64	60	71	58	2,880,000	IDAHO							
Little Tennessee Project: Nantahala, Thorpe, Fontana, and Chilhowee Lakes (FPR)	55	60	84	52	1,478,000	Boise River (4 reservoirs) (FIP)	81	93	82	77	1,235,000		
WISCONSIN													
Chippewa and Flambeau (PR)	91	96	86	94	365,000	Coeur d'Alene Lake (P)	96	121	125	70	238,500		
Wisconsin River (21 reservoirs) (PR)	79	91	82	87	399,000	Pend Oreille Lake (FP)	48	85	82	51	1,561,000		
MINNESOTA													
Mississippi River headwater system (FMR)	53	41	37	44	1,640,000	IDAHO-WYOMING							
NORTH DAKOTA													
Lake Sakakawea (Garrison) (FIPR)	84	84	86	79	22,700,000	Upper Snake River (8 reservoirs) (MP)	62	85	79	59	4,401,000		
SOUTH DAKOTA													
Angostura (I)	94	74	90	92	127,600	WYOMING							
Belle Fourche (I)	86	77	74	62	185,200	Boysen (FIP)	64	70	67	62	802,000		
Lake Francis Case (FIP)	87	76	82	86	4,834,000	Buffalo Bill (IP)	70	68	74	70	421,300		
Lake Oahe (FIP)	100	88	100	97	22,530,000	Keyhole (F)	39	40	50	35	193,800		
Lake Sharpe (FIP)	101	100	100	101	1,725,000	Pathfinder, Seminole, Alcona, Kortes, Glendo, and Guernsey Reservoirs (I)	78	83	61	72	3,056,000		
Lewis and Clark Lake (FIP)	76	77	85	77	477,000	COLORADO							
ARIZONA-NEVADA													
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	John Martin (FIR)	59	97	17	71	364,400		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	Taylor Park (IR)	59	84	70	46	106,200		
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Colorado-Big Thompson project (I)	75	87	64	93	730,300		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	COLORADO RIVER STORAGE PROJECT							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Lake Powell; Flaming Gorge, Fontenelle, Navajo, and Blue Mesa Reservoirs (IFPR)	87	94	71	85	31,620,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	UTAH-IDAHO							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Bear Lake (IPR)	95	89	70	86	1,421,000		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	CALIFORNIA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Folsom (FIP)	88	88	88	69	1,000,000		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	Hetch Hetchy (MP)	68	82	70	71	360,400		
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Isabella (FIR)	89	66	48	79	568,100		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	Pine Flat (FI)	82	84	73	73	1,001,000		
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Clair Engle Lake (Lewiston) (P)	96	87	91	92	2,438,000		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	Lake Almanor (P)	106	74	66	100	1,036,000		
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Lake Berryessa (FIMW)	98	87	88	100	1,600,000		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	Millerton Lake (FI)	61	84	78	42	503,200		
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Shasta Lake (FIPR)	93	54	91	93	4,377,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	CALIFORNIA-NEVADA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Lake Tahoe (IPR)	93	82	68	88	744,600		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	NEVADA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Rye Patch (I)	95	91	72	94	194,300		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	ARIZONA-NEVADA							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	ARIZONA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	San Carlos (IP)	90	99	27	95	935,100		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	Salt and Verde River system (IMPR)	94	98	54	98	2,019,100		
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	NEW MEXICO							
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	Conchas (FIR)	78	85	79	81	330,100		
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	NEVADA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Rye Patch (I)	95	91	72	94	194,300		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	ARIZONA-NEVADA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	ARIZONA							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	San Carlos (IP)	90	99	27	95	935,100		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	Salt and Verde River system (IMPR)	94	98	54	98	2,019,100		
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	NEW MEXICO							
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	Conchas (FIR)	78	85	79	81	330,100		
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	NEVADA							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Rye Patch (I)	95	91	72	94	194,300		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	ARIZONA-NEVADA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	ARIZONA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	San Carlos (IP)	90	99	27	95	935,100		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	Salt and Verde River system (IMPR)	94	98	54	98	2,019,100		
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	NEW MEXICO							
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	Conchas (FIR)	78	85	79	81	330,100		
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	NEVADA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Rye Patch (I)	95	91	72	94	194,300		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	ARIZONA-NEVADA							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	ARIZONA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	San Carlos (IP)	90	99	27	95	935,100		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	Salt and Verde River system (IMPR)	94	98	54	98	2,019,100		
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	NEW MEXICO							
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	Conchas (FIR)	78	85	79	81	330,100		
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	NEVADA							
ARIZONA													
San Carlos (IP)	90	99	27	95	935,100	Rye Patch (I)	95	91	72	94	194,300		
Salt and Verde River system (IMPR)	94	98	54	98	2,019,100	ARIZONA-NEVADA							
NEW MEXICO													
Conchas (FIR)	78	85	79	81	330,100	Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000		
Elephant Butte and Caballo (FIPR)	93	89	34	95	2,453,000	ARIZONA							
NEVADA													
Rye Patch (I)	95	91	72	94	194,300	San Carlos (IP)	90	99	27	95	935,100		
ARIZONA-NEVADA													
Lake Mead and Lake Mohave (FIMP)	92	93	71	90	27,970,000	Salt and Verde River system (IMPR)	94	98	54	98	2,019,100		

¹ acre-foot = 0.04356 million cubic feet = 0.326 million gallons = 0.504 cubic feet per second day.
Thousands of kilowatt-hours (the potential electric power that could be generated by the volume of water in storage).

USABLE CONTENTS OF SELECTED RESERVOIRS AND RESERVOIR SYSTEMS



FLOW OF LARGE RIVERS DURING MAY 1986

Station number	Stream and place of determination	Drainage area (square miles)	Average discharge through September 1980 (cubic feet per second)	May 1986					
				Monthly mean discharge (cubic feet per second)	Percent of median monthly discharge, 1951-80	Change in discharge from previous month (percent)	Discharge near end of month		
							Cubic feet per second	Million gallons per day	Date
01014000	St. John River below Fish River at Fort Kent, ME.	5,690	9,647	17,000	50	-49	14,900	9,630	31
01318500	Hudson River at Hadley, NY.....	1,664	2,909	3,750	75	-54	3,950	2,552	31
01357500	Mohawk River at Cohoes, NY.....	3,456	5,734	5,220	78	-46	2,500	1,620	31
01463500	Delaware River at Trenton, NJ.....	6,780	11,750	12,650	100	-29	9,860	6,372	31
01570500	Susquehanna River at Harrisburg, PA.	24,100	34,530	24,160	57	-53	24,700	15,960	29
01646500	Potomac River near Washington, DC.	11,560	¹ 11,490	¹ 7,580	54	-42	5,560	3,593	31
02105500	Cape Fear River at William O. Huske Lock near Tarheel, NC.	4,810	5,005	1,180	35	-19	1,200	780	31
02131000	Pee Dee River at Peedee, SC.....	8,830	9,851	2,516	33	-29	3,530	2,281	29
02226000	Altamaha River at Doctortown, GA....	13,600	13,880	2,960	24	-60	2,580	1,667	29
02320500	Suwannee River at Branford, FL.....	7,880	6,987	5,090	77	-47	4,140	2,675	31
02358000	Apalachicola River at Chattahoochee, FL.	17,200	22,570	9,190	46	-23	8,560	5,532	31
02467000	Tombigbee River at Demopolis lock and dam near Coatopa, AL.	15,400	23,300	6,881	31	+50	32,000	20,700	30
02489500	Pearl River near Bogalusa, LA.....	6,630	9,768	5,163	50	+32	32,000	20,700	31
03049500	Allegheny River at Natrona, PA.....	11,410	¹ 19,480	¹ 10,180	48	-54	9,000	5,800	27
03085000	Monongahela River at Braddock, PA..	7,337	¹ 12,510	¹ 8,244	59	-27	7,400	4,780	21
03193000	Kanawha River at Kanawha Falls, WV.	8,367	12,590	9,646	75	+90	8,710	5,629	27
03234500	Scioto River at Higby, OH.....	5,131	4,547	1,597	34	-29	1,920	1,240	30
03294500	Ohio River at Louisville, KY ²	91,170	116,000	64,120	49	-14	58,730	37,960	26
03377500	Wabash River at Mount Carmel, IL....	28,635	27,220	30,690	96	+36	39,100	25,270	30
03469000	French Broad River below Douglas Dam, TN.	4,543	6,798	3,085	46	+2
04084500	Fox River at Rapide Croche Dam, near Wrightstown, WI. ²	6,150	4,163	4,463	78	-59	2,843	1,837	31
04264331	St. Lawrence River at Cornwall, Ontario-near Massena, NY. ³	299,000	242,700	331,300	119	+4	319,000	206,200	31
02NG001	St. Maurice River at Grand Mere, PQ.	16,300	25,150	70,700	103	+18	39,100	25,270	30
05082500	Red River of the North at Grand Forks, ND.	30,100	2,551	14,740	390	-23	14,200	9,180	21
05133500	Rainy River at Manitou Rapids, MN...	19,400	11,830	33,800	187	+69	31,500	20,360	27
05330000	Minnesota River near Jordan, MN.....	16,200	3,402	22,250	404	-19	14,300	9,240	31
05331000	Mississippi River at St. Paul, MN.....	36,800	¹ 10,610	64,230	290	-6	39,000	25,200	31
05365500	Chippewa River at Chippewa Falls, WI.	5,600	5,100	6,133	101	-65	2,000	1,300	31
05407000	Wisconsin River at Muscoda, WI.....	10,300	8,617	8,145	73	-62	6,484	4,190	30
05446500	Rock River near Joslin, IL.....	9,551	5,873	10,700	157	-16	10,000	6,000	31
05474500	Mississippi River at Keokuk, IA.....	119,000	62,620	178,500	184	-7	209,000	135,100	31
06214500	Yellowstone River at Billings, MT....	11,796	7,038	11,300	85	+131	29,900	19,320	30
06934500	Missouri River at Hermann, MO.....	524,200	79,490	144,000	156	+33	147,000	95,000	30
07289000	Mississippi River at Vicksburg, MS ⁴ ...	1,140,500	576,600	592,400	71	-9	764,000	493,800	27
07331000	Washita River near Dickson, OK.....	7,202	1,368	3,760	225	+164	3,000	1,900	21
08276500	Rio Grande below Taos Junction Bridge, near Taos, NM.	9,730	725	2,103	237	+44	2,300	1,490	30
09315000	Green River at Green River, UT.....	40,600	6,298	21,020	182	+58
11425500	Sacramento River at Verona, CA.....	21,257	18,820	11,002	62	-44	12,700	8,210	28
13269000	Snake River at Weiser, ID.....	69,200	18,050	41,500	159	-15	43,900	28,370	31
13317000	Salmon River at White Bird, ID.....	13,550	11,250	27,600	87	+49	101,100	65,340	31
13342500	Clearwater River at Spalding, ID.....	9,570	15,480	31,800	63	+48	65,600	42,400	31
14105700	Columbia River at The Dalles, OR ⁵	237,000	¹ 193,100	¹ 313,600	73	+25	268,100	173,300	27
14191000	Willamette River at Salem, OR.....	7,280	¹ 23,510	¹ 21,500	92	+24	14,500	9,370	27
15515500	Tanana River at Nenana, AK.....	25,600	23,460	27,920	95	+278	35,000	22,600	31
08MF005	Fraser River at Hope, BC.....	83,800	96,290	130,300	72	+79	269,800	174,400	30

¹Adjusted.²Records furnished by Corps of Engineers.³Records furnished by Buffalo District, Corps of Engineers, through International St. Lawrence River Board of Control. Discharges shown are considered to be the same as discharge at Ogdensburg, N.Y. when adjusted for storage in Lake St. Lawrence.⁴Records of daily discharge computed jointly by Corps of Engineers and Geological Survey.⁵Discharge determined from information furnished by Bureau of Reclamation, Corps of Engineers, and Geological Survey.

DISSOLVED SOLIDS AND WATER TEMPERATURES, FOR MAY 1986, AT DOWNSTREAM SITES ON SIX LARGE RIVERS

Station number	Station name	May data of following calendar years	Stream discharge during month	Dissolved-solids concentration ^a		Dissolved-solids discharge ^a			Water temperature ^b		
			Mean (cfs)	Mini-mum (mg/L)	Maxi-mum (mg/L)	Mean	Mini-mum	Maxi-mum	Mean in °C	Mini-mum, in °C	Maxi-mum, in °C
						(tons per day)					
01463500	Delaware River at Trenton, NJ (Morrisville, PA).	1986 1945—85 (Extreme yr)	*11,410 15,370 °12,650	84 50 (1946)	121 123 (1978)	2,590 (1965)	1,880 930 (1965)	3,810 21,790 (1984)	17.0 ... 10.0	14.0 10.0	21.0 28.5
04264331	St. Lawrence River at Cornwall, Ontario, near Massena, NY (median streamflow at Ogdensburg, NY).	1986 1976—85 (Extreme yr)	331,300 293,300 °278,300	166 165 (**)	167 167 (**)	148,900 131,600	142,900 109,000 (1981)	153,400 153,000 (1976)	10.0 9.0	7.0 4.0	12.5 13.0
07289000	Mississippi River at Vicksburg, MS.	1986 1976—85 (Extreme yr)	592,400 895,500 °838,200	240 178 (1977)	281 290 (1982)	411,000 511,500	347,000 176,000 (1977)	561,000 954,000 (1983)	23.0 20.0	19.5 14.5	25.5 26.0
03612500	Ohio River at lock and dam 53, near Grand Chain, IL (streamflow station at Metropolis, IL).	1986 1955—85 (Extreme yr)	132,000 365,900 °296,000	173 124 (1983)	235 287 (1979)	47,300 25,500 (1976)	95,300 466,000 (1984) 16.0	16.0 6.5	22.0 25.0
06934500	Missouri River at Hermann, MO (60 miles west of St. Louis, MO).	1986 1976—85 (Extreme yr)	164,000 123,500 °92,040	285 211 (1978)	468 520 (1981)	157,000 113,200	100,000 44,500 (1977)	237,000 272,000 (1983)	21.0 18.5	19.0 13.0	22.5 24.5
14128910	Columbia River at Warrendale, OR (streamflow station at The Dalles, OR).	1986 1976—85 (Extreme yr)	249,000 263,400 °427,700	90 67 (1976)	97 144 (1977)	63,200 68,100	48,000 37,500 (1977)	77,300 102,000 (1983)	11.5 12.5	9.5 9.5	14.5 16.5

^aDissolved-solids concentrations, when not analyzed directly, are calculated on basis of measurements of specific conductance.


^bTo convert °C to °F: [(1.8 X °C) + 32] = °F.

^cMedian of monthly values for 30-year reference period, water years 1951—80, for comparison with data for current month.

^dDissolved solids and water temperature records are for the first 18 days of the month.

**Occurred several years

PRECIPITATION FORECAST FOR JUNE THROUGH AUGUST 1986





(From Monthly and Seasonal Weather Outlook Published by National Weather Service)

STREAMFLOW DURING MAY



1981

1982





1983


1984




STREAMFLOW

 Above-normal
(with the highest 25 percent
of record for this month)

 In normal range

 Below normal
(within the lowest 25 percent
of record for the month)

MAY 1985



NATIONAL WATER CONDITIONS

MAY 1986

Based on reports from the Canadian and U.S. Field offices; completed July 9, 1986

TECHNICAL STAFF

Thomas G. Ross, Editor
Carroll W. Saboe
John C. Kammerer
Allen Sinnott
Krishnaveni V. Sarma
Sharon A. Edmonds
Carole J. Marlow

COPY PREPARATION

Lois C. Fleshmon
Sharon L. Peterson
Aisha P.R. Law

GRAPHICS

Frances B. Davison
Carolyn L. Moss

The National Water Conditions is published monthly. Subscriptions are free on application to the U.S. Geological Survey, 419 National Center, Reston, VA 22092.

EXPLANATION OF DATA (Revised January 1986)

Cover map shows generalized pattern of streamflow for the month based on provisional data from 184 index gaging stations—18 in Canada, 164 in the United States, and 2 in the Commonwealth of Puerto Rico. Alaska and Hawaii inset maps show streamflow only at the index gaging stations that are located near the point shown by the arrows. Classifications on map are based on comparison of streamflow for the current month at each index station with the flow for the same month in the 30-year reference period, 1951–80. Shorter reference periods are used for one Canadian index station, two Kansas index stations, one New York index station, and the Puerto Rico index stations because of the limited records available.

The comparative data are obtained by ranking the 30 flows for each month of the reference period in order of decreasing magnitude—the highest flow is given a ranking of 1 and the lowest flow is given a ranking of 30. Quartiles (25-percent points) are computed by averaging the 7th and 8th highest flows (upper quartile), 15th and 16th highest flows (middle quartile and median), and the 23rd and 24th highest flows (lower quartile). The upper and lower quartiles set off the highest 25 percent

of flows and lowest 25 percent of flows, respectively, for the reference period. The median (middle quartile) is the middle value by definition. For the reference period, 50 percent of the flows are greater than the median, 50 percent are less than the median, 50 percent are between the upper and lower quartiles (in the normal range) 25 percent are greater than the upper quartile (above normal), and 25 percent are less than the lower quartile (below normal). Flow for the current month is then classified as; *above normal* if it is greater than the upper quartile, *in the normal range* if it is between the upper and lower quartiles, and *below normal* if it is less than the lower quartile. Change in flow from the previous month to the current month is classified as *seasonal* if the change is in the same direction as the change in the median. If the change is in the opposite direction of the change in the median, the change is classified as *contraseasonal* (opposite to the seasonal change). For example: at a particular index station, the January median is greater than the December median; if flow for the current January increased from December (the previous month), the increase is seasonal; if flow for the current January decreased from December, the decrease is *contraseasonal*.

Flood frequency analyses define the relation of flood peak magnitude to probability of occurrence or recurrence interval. *Probability of occurrence* is the chance that a given flood magnitude will be exceeded in any one year. *Recurrence interval* is the reciprocal of probability of occurrence and is the average number of years between occurrences. For example, a flood having a probability of occurrence of 0.01 (1 percent) has a recurrence interval of 100 years. *Recurrence intervals imply no regularity of occurrence*; a 100-year flood might be exceeded in consecutive years or it might not be exceeded in a 100-year period.

Statements about *ground-water levels* refer to conditions near the end of the month. The water level in each key observation well is compared with average level for the end of the month determined from the 30-year reference period, 1951–80, or from the entire past record for that well when only limited records are available. Comparative data for ground-water levels are obtained in the same manner as comparative data for streamflow. *Changes in ground-water levels*, unless described otherwise, are from the end of the previous month to the end of the current month.

Dissolved solids and temperature data for May are given for six stream-sampling sites that are part of the National Stream Quality Accounting Network (NASQAN). *Dissolved solids* are minerals dissolved in water and usually consist predominately of silica and ions of calcium, magnesium, sodium, potassium, carbonate, bicarbonate, sulfate, chloride, and nitrate. *Dissolved-solids discharge* represents the total daily amount of dissolved minerals carried by the stream. *Dissolved-solids concentrations* are generally higher during periods of low streamflow, but the highest dissolved-solids discharges occur during periods of high streamflow because the total quantities of water, and therefore total load of dissolved minerals, are so much greater than at times of low flow.

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY
NATIONAL CENTER, STOP 419
RESTON, VIRGINIA 22092

OFFICIAL BUSINESS

Return this sheet to above address, if you do
NOT wish to receive this material ☐, or
if change of address is needed ☐ (indicate
change, including ZIP code).

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF THE INTERIOR
INT 413



FIRST CLASS

SPECIAL PROCESSING DEPT
MARCIA KOZLOWSKI
XEROX/UNIVERSITY MICROFILMS
ANN ARBOR MI 48106

NAC

